

ORBIFOLD QUANTUM COHOMOLOGY OF THE CLASSIFYING SPACE OF A FINITE GROUP

TYLER J. JARVIS AND TAKASHI KIMURA

ABSTRACT. We work through, in detail, the quantum cohomology, with gravitational descendants, of the orbifold $\mathcal{B}G$, the point with action of a finite group G . We provide a simple description of algebraic structures on the state space of this theory. As a consequence, we find that multiple copies of commuting Virasoro algebras appear which completely determine the correlators of the theory.

1. INTRODUCTION

W. Chen and Y. Ruan [CR1, CR2] introduced the notion of the Gromov-Witten invariants of an orbifold V . Their construction reduces to the usual Gromov-Witten invariants when V is a smooth, projective variety. When $V = [Y/G]$, where Y is a smooth, projective variety, and G is a finite group, the state space of this theory is generally larger than the G -invariant part of the cohomology of Y ; indeed, it has additional direct summands associated to loci in Y with nontrivial isotropy. These loci are called the twisted sectors of the theory, and their presence should be part of the proper notion of the cohomology of an orbifold.

The purpose of this paper is to provide a detailed treatment of this theory for the simplest case, namely, when V is the classifying stack $\mathcal{B}G$ of a finite group G . Many of the features and subtleties of the Gromov-Witten invariants of orbifolds are present even here. The state space of this theory contains twisted sectors and the correlators in this theory are intersection numbers on $\overline{\mathcal{M}}_{g,n}(\mathcal{B}G)$, the moduli space of genus- g , n -pointed orbifold stable maps into $\mathcal{B}G$.

The correlators in this theory can be described in purely group theoretic terms, and we recover the result [Ru] that the algebraic structure on the state space $\mathcal{H}$ is isomorphic to the center of the group algebra, $Z\mathbb{C}[G]$, together with an invariant metric.

Furthermore, we show that on the large phase space of this theory, there are r commuting copies of “half” the Virasoro algebra in this theory, where r is the dimension of $\mathcal{H}$, all of which annihilate, and completely determine, the exponential of the large phase space potential function. We obtain a proof of the usual Virasoro conjecture [EHX] as the special case where a diagonal action (after a variable rescaling) of these Virasoro algebras is considered. Similarly, the relevant integrable hierarchy consists of r commuting copies of the KdV hierarchy.

Finally, it is worth pointing out that the moduli spaces $\overline{\mathcal{M}}_{g,n}(\mathcal{B}G)$ have many features similar to $\overline{\mathcal{M}}_{g,n}^{1/r}$, the moduli space of r -spin curves [JKV]. The moduli spaces $\overline{\mathcal{M}}_{g,n}(\mathcal{B}G)$ have boundary strata indexed by stable graphs whose tails and

2000 *Mathematics Subject Classification.* Primary: 14N35, 53D45. Secondary: 20C05, 14H10. Research of the first author was partially supported by NSF grant DMS-0105788.

half-edges are decorated by elements of G (up to conjugation) while $\overline{\mathcal{M}}_{g,n}^{1/r}$ have boundary strata indexed by stable graphs whose tails and half-edges are decorated by elements of $\mathbb{Z}_r$. The construction of the correlators in both theories is also analogous. However, the ring structures on the state spaces are distinctly different. It would be interesting to find the analogs of the above results for the spaces $\overline{\mathcal{M}}_{g,n}^{1/r}$.

1.1. Conventions and notation. Throughout this paper, except where otherwise specified, we will work only over the complex numbers $\mathbb{C}$, and all groups will be finite.

For a given group G we denote the classifying stack of G by $\mathcal{B}G$; namely, $\mathcal{B}G$ is the stack quotient

$$\mathcal{B}G := [pt/G]$$

of a point modulo a trivial G action.

Group elements will always be denoted by lower-case Greek letters, and the conjugacy class of an element $\gamma \in G$ is denoted $[\gamma]$. An n -tuple of elements $(\gamma_1, \dots, \gamma_n)$ will generally be denoted by a boldfaced γ . The centralizer in G of an element γ is denoted $C(\gamma)$, and the intersection $\bigcap_{i=1}^n C(\gamma_i)$ of several centralizers is denoted $C(\gamma_1, \dots, \gamma_n) = C(\gamma)$. The commutator $\alpha\beta\alpha^{-1}\beta^{-1}$ of two group elements α and β is denoted $[\alpha, \beta]$. Finally, the center of an algebra A will be denoted ZA .

Acknowledgments We would like to thank Dan Abramovich and Yongbin Ruan for many helpful discussions, and Heidi Jarvis for help with typesetting.

2. ORBIFOLD STABLE MAPS INTO $\mathcal{B}G$

2.1. The stack. Our chief objects of study in this paper are Gromov-Witten invariants arising from the stack $\overline{\mathcal{M}}_{g,n}(\mathcal{B}G)$ of n -pointed orbifold stable maps into $\mathcal{B}G$, in the sense of W. Chen and Y. Ruan [CR1, CR2] (called balanced twisted stable maps by Abramovich and Vistoli [AV]).

These are maps $f : \Sigma \longrightarrow \mathcal{B}G$ from an orbifold Riemann surface (orbicurve) Σ into the classifying stack $\mathcal{B}G$ of a finite group G . Here Σ has non-trivial orbifold structure only at marked points $p_1, \dots, p_n$ and at nodes, and the orbifold structure at the nodes is *balanced*, meaning that the action of the stabilizer $G_q \simeq \mu_l$ at a nodal geometric point q of Σ has complementary eigenvalues on the tangent spaces of the two branches of Σ at q .

The stack $\overline{\mathcal{M}}_{g,n}(\mathcal{B}G)$ is a smooth, proper Deligne-Mumford stack with projective coarse moduli space [ACV, Thm 3.0.2]. This stack has a number of important connections to other moduli problems. For example, in the case that G is the symmetric group S_d on d letters, the stack $\overline{\mathcal{M}}_{g,n}(\mathcal{B}S_d)$ is the normalization of the stack of admissible covers [ACV, §4].

Recall that an orbifold stable map from a smooth, n -pointed orbicurve $(\Sigma, p_1, \dots, p_n)$ into $\mathcal{B}G$ determines a principal G bundle on the complement $\Sigma - \{p_1, \dots, p_n\}$. This, in turn, is determined by its holonomy, that is by a homomorphism $\pi_1(\Sigma - \{p_1, \dots, p_n\}) \longrightarrow G$. Moreover, the stabilizer G_{p_i} of a marked point p_i of Σ is always cyclic, and the order of G_{p_i} is equal to the order of the holonomy around that marked point. Conversely, G acts by conjugation on the homomorphisms $\pi_1(\Sigma - \{p_1, \dots, p_n\}) \longrightarrow G$, and two such homomorphisms determine the same G -bundle on $\Sigma - \{p_1, \dots, p_n\}$ precisely when they differ by this adjoint action of G .

Finally, since $\mathcal{BG}$ is a proper, separated stack, any principal G bundle on $\Sigma - \{p_1, \dots, p_n\}$ extends uniquely, after suitable base extension, to a principal bundle on some proper curve $\tilde{\Sigma}$, with an isomorphism $\tilde{\Sigma} - \{p_1, \dots, p_n\} \longrightarrow \Sigma - \{p_1, \dots, p_n\}$. The data of this cover (up to the obvious notion of equivalence of such data) is exactly equivalent to the data of a principal bundle on an orbicurve. Moreover, the adjoint action of G on these homomorphisms exactly corresponds to the natural action of G on the orbifold stable maps. Thus we have the following:

Proposition 2.1. *For a given curve $[C, p_1, \dots, p_n]$ corresponding to a point of the smooth locus $\mathcal{M}_{g,n}$, the fiber of $\overline{\mathcal{M}}_{g,n}(\mathcal{BG})$ over the point $[C]$ corresponds to the quotient $\text{Hom}(\pi_1(C - \{p_1, \dots, p_n\}), G)/\text{ad } G$.*

2.2. Morphisms. There are several natural morphisms from $\overline{\mathcal{M}}_{g,n}(\mathcal{BG})$.

First, recall that for $X = \mathcal{BG}$ the twisted sectors $X_{[\gamma]} = \{(x, [\gamma]) | x \in X, \gamma \in G_x\}$ of [CR1] are simply $\mathcal{BG}_{[\gamma]} \cong [pt/C(\gamma)]$.

There are evaluation morphisms

$$ev_i : \overline{\mathcal{M}}_{g,n}(\mathcal{BG}) \longrightarrow \widetilde{\mathcal{BG}},$$

where $\widetilde{\mathcal{BG}} = \coprod_{[\gamma]} \mathcal{BG}_{[\gamma]}$ is the disjoint union of twisted sectors. We can describe the evaluation morphism ev_i as follows: Any stable map $f : \Sigma \longrightarrow \mathcal{BG}$ must be representable, and hence must induce an injective homomorphism $f_* : G_{p_i} \longrightarrow G$ from the local group $G_{p_i} = \mathbb{Z}/m_i$ of the i th marked point p_i of Σ into G . The group G acts by conjugation on this homomorphism f_* , and so the image of $1 \in \mathbb{Z}/m_i$ in G is defined only up to conjugacy. The evaluation morphism ev_i is the morphism taking $[\Sigma, p_1, \dots, p_n, f]$ to the point $(f(p_i), [f_*(1)]) \in \mathcal{BG}_{[f_*(1)]} \subseteq \widetilde{\mathcal{BG}}$. Alternatively, the image $f_*(1)$ is simply the holonomy of the induced G -bundle on $\Sigma - \{p_1, \dots, p_n\}$ around the marked point p_i .

We can use the evaluation morphism to see that the stack $\overline{\mathcal{M}}_{g,n}(\mathcal{BG})$ breaks up as the disjoint union of open and closed substacks

$$\overline{\mathcal{M}}_{g,n}(\mathcal{BG}) = \coprod_{([\gamma_1], \dots, [\gamma_n])} \overline{\mathcal{M}}_{g,n}(\mathcal{BG}, [\gamma_1], \dots, [\gamma_n]),$$

where $\overline{\mathcal{M}}_{g,n}(\mathcal{BG}, [\gamma_1], \dots, [\gamma_n]) = ev_1^{-1}(\mathcal{BG}_{[\gamma_1]}) \cap \dots \cap ev_n^{-1}(\mathcal{BG}_{[\gamma_n]})$ is the substack of points of $\overline{\mathcal{M}}_{g,n}(\mathcal{BG})$ mapped by ev_i to $\mathcal{BG}_{\gamma_i}$ for every $i \in \{1, \dots, n\}$. Of course, $\overline{\mathcal{M}}_{g,n}(\mathcal{BG}, [\gamma_1], \dots, [\gamma_n])$ may be empty for some choices of conjugacy classes $([\gamma_1], \dots, [\gamma_n])$.

In the special case of $\overline{\mathcal{M}}_{0,3}(\mathcal{BG})$, since there is only one 3-pointed, genus-zero stable curve (call it Σ), we may fix, once and for all a base point q (distinct from the three marked points p_1, p_2 , and p_3) and a basis $\{s_1, s_2, s_3\}$ of $\pi_1(\Sigma - \{p_1, p_2, p_3\}, q)$ such that $\prod_{i=1}^3 s_i = 1$. In this case, Proposition 2.1 shows that each component of $\overline{\mathcal{M}}_{0,3}(\mathcal{BG})$ is uniquely determined up to simultaneous (diagonal) adjoint action of G by a choice of three elements $\gamma_1, \gamma_2, \gamma_3 \in G$ such that $\prod \gamma_i = 1$. That is, if we let $[\gamma]$ denote the diagonal conjugacy class of the triple $\gamma = (\gamma_1, \gamma_2, \gamma_3)$, and we let T^3 denotes the set of all such triple conjugacy classes, whose product is trivial (i.e., $\prod \gamma_i = 1$), then

$$\overline{\mathcal{M}}_{0,3}(\mathcal{BG}) = \coprod_{[\gamma] \in T^3} \overline{\mathcal{M}}_{0,3}(\mathcal{BG}, [\gamma]),$$

and for a given $[\gamma] \in T^3$, we have

$$\overline{\mathcal{M}}_{0,3}(\mathcal{B}G, [\gamma]) \cong \mathcal{B}C(\gamma).$$

For $g > 0$ or $n > 3$ one cannot index the moduli space so easily, since deformations in the moduli space may act on the holonomies in non-diagonal ways. Nevertheless, for any fixed, smooth, n -pointed curve, Proposition 2.1 gives a complete description of the points of $\overline{\mathcal{M}}_{g,n}(\mathcal{B}G)$ that lie over it. This shows that the forgetful morphism $\pi : \overline{\mathcal{M}}_{g,n}(\mathcal{B}G) \longrightarrow \overline{\mathcal{M}}_{g,n}$, is quasi-finite. In fact π is also proper, but it is not generally representable [ACV, Cor 3.0.5].

Finally we have the forgetting-tails morphism, defined as follows. When the holonomy γ_i around a marked point p_i is trivial, then we may *forget* the data of that marked point. This gives a morphism

$$\overline{\mathcal{M}}_{g,n+1}(\mathcal{B}G, [\gamma_1], \dots, [1], \dots, [\gamma_n]) \longrightarrow \overline{\mathcal{M}}_{g,n}(\mathcal{B}G, [\gamma_1], \dots, [\widehat{1}], \dots, [\gamma_n]).$$

Note that the forgetting-tails morphism is not defined for all of $\overline{\mathcal{M}}_{g,n+1}(\mathcal{B}G)$, but rather only for the components corresponding to marked points with trivial holonomy.

3. GROMOV-WITTEN INVARIANTS, COHOMOLOGICAL FIELD THEORY AND K-THEORY

We define the classes ψ_i on $\overline{\mathcal{M}}_{g,n}(\mathcal{B}G)$ to be the pullbacks $\psi_i = \pi^*(\psi_i)$ of the ψ_i classes on $\overline{\mathcal{M}}_{g,n}$.

The tangent bundle of $\mathcal{B}G$ is trivial, and thus the virtual fundamental class of $\overline{\mathcal{M}}_{g,n}(\mathcal{B}G)$ is just the usual fundamental class.

The orbifold cohomology of $\mathcal{B}G$ is, as a vector space,

$$\mathcal{H} := H_{orb}^*(\mathcal{B}G, \mathbb{C}) := H^*(\widetilde{\mathcal{B}G}, \mathbb{C}) = \bigoplus_{[\gamma]} \mathbb{C}.$$

For each conjugacy class $[\gamma]$ in G , let $e_{[\gamma]}$ denote the class $1 \in H^0(\mathcal{B}G_{[\gamma]}, \mathbb{C}) \subseteq \mathcal{H}$. The $e_{[\gamma]}$'s form a basis of $\mathcal{H}$ and we may form n -point correlators

$$\langle \tau_{a_1}(e_{[\gamma_1]}) \dots \tau_{a_n}(e_{[\gamma_n]}) \rangle_g^G := \int_{\overline{\mathcal{M}}_{g,n}(\mathcal{B}G)} \prod_i \psi_i^{a_i} ev_i^*(e_{[\gamma_i]}).$$

3.1. Cohomological field theory. The three-point, genus-zero correlators play a special role, since they define the metric on $\mathcal{H}$ and the quantum (orbifold) product. They vanish for dimensional reasons unless $\sum_{i=1}^n a_i = 0$.

Proposition 3.1. *We have*

$$\begin{aligned} \langle \tau_0(e_{[\gamma_1]}) \tau_0(e_{[\gamma_2]}) \tau_0(e_{[\gamma_3]}) \rangle_0^G &= \sum_{\substack{\sigma_1, \sigma_2, \sigma_3 \\ \prod \sigma_i = 1 \\ \sigma_i \in [\gamma_i]}} \frac{1}{|G|} \\ &= \sum_{\substack{[\sigma_1, \sigma_2, \sigma_3] \\ \prod \sigma_i = 1 \\ \sigma_i \in [\gamma_i]}} \frac{1}{|C(\sigma_1, \sigma_2)|}. \end{aligned}$$

Proof. For a given component $\overline{\mathcal{M}}_{0,3}(\mathcal{B}G, [\sigma_1, \sigma_2, \sigma_3])$ of the moduli space $\overline{\mathcal{M}}_{0,3}(\mathcal{B}G)$, the evaluation map ev_i maps this component to $\mathcal{B}G_{[\sigma_i]}$, so

$$ev_i^*(e_{[\gamma_i]}) = \begin{cases} 0 & \text{if } \sigma_i \notin [\gamma_i] \\ 1 & \text{if } \sigma_i \in [\gamma_i] \end{cases}.$$

Moreover,

$$\int_{\overline{\mathcal{M}}_{0,3}(\mathcal{B}G, [\sigma_1, \sigma_2, \sigma_3])} 1 = \deg(\overline{\mathcal{M}}_{0,3}(\mathcal{B}G, [\sigma_1, \sigma_2, \sigma_3])) = \frac{1}{|C(\sigma_1, \sigma_2, \sigma_3)|} = \frac{1}{|C(\sigma_1, \sigma_2)|}.$$

So

$$(3.1) \quad \langle \tau_0(e_{[\sigma_1]})\tau_0(e_{[\sigma_2]})\tau_0(e_{[\sigma_3]}) \rangle_0^G = \sum_{\substack{[\sigma_1, \sigma_2, \sigma_3] \\ \prod_{i=1}^3 \sigma_i = 1 \\ \sigma_i \in [\gamma_i]}} \frac{1}{|C(\sigma_1, \sigma_2)|},$$

and it is easy to see that the number of elements in a given non-empty conjugacy class $[\sigma_1, \sigma_2, \sigma_3]$ of triples is exactly $\frac{|G|}{|C(\sigma_1, \sigma_2)|}$, so the expression (3.1) gives the rest of the proposition. $\square$

Corollary 3.2. *The metric on $\mathcal{H}$ induced by the 3-point correlators is*

$$\eta_{[\gamma_1][\gamma_2]} := \eta(e_{[\gamma_1]}, e_{[\gamma_2]}) = \frac{1}{|C(\gamma_1)|} \delta_{[\gamma_1][\gamma_2^{-1}]},$$

which is non-degenerate on $\mathcal{H}$.

The quantum product is given by

$$e_{[\gamma_1]} * e_{[\gamma_2]} = \sum_{\substack{\sigma_1, \sigma_2 \\ \sigma_i \in [\gamma_i]}} \frac{|C(\sigma_1, \sigma_2)|}{|G|} e_{[\sigma_1, \sigma_2]}.$$

It is clear that $\mathcal{H}$ is additively isomorphic to the ring $\text{Class}_{\mathbb{C}}(G)$ of class functions of G , where $e_{[\gamma]}$ is the class function $e_{[\gamma]}(\sigma) = \delta_{[\gamma], [\sigma]}$. Moreover, $\text{Class}_{\mathbb{C}}(G)$ has a natural metric on it: $\langle f, g \rangle = \frac{1}{|G|} \sum_{\sigma \in G} f(\sigma)g(\sigma^{-1})$. Let Φ denote the standard additive isomorphism

$$\Phi : \text{Class}_{\mathbb{C}}(G) \longrightarrow Z\mathbb{C}[G]$$

defined by linearly extending $\Phi(e_{[\gamma]}) = \sum_{\alpha \in [\gamma]} \alpha$. The standard group-algebra product $\cdot$ in $Z\mathbb{C}[G]$ pulls back via Φ to convolution $\star$ of functions. And the push-forward $\Phi_* \langle \cdot, \cdot \rangle$ to $Z\mathbb{C}[G]$ of the metric on $\text{Class}_{\mathbb{C}}(G)$ agrees with the metric $(\cdot, \cdot)$ on $\mathbb{C}[G]$ defined as $(\alpha, \beta) = \frac{1}{|G|} \delta_{\alpha, \beta^{-1}}$, when α and β are in G . With respect to this metric, $Z\mathbb{C}[G]$ is a Frobenius algebra.

Corollary 3.3. *The homomorphisms*

$$(\mathcal{H}, *, \eta) \longrightarrow (\text{Class}_{\mathbb{C}}(G), \star, \langle \cdot, \cdot \rangle) \xrightarrow{\Phi} (Z\mathbb{C}[G], \cdot, (\cdot, \cdot))$$

are isomorphisms of Frobenius algebras.

Proof. It is well-known, and clear, that these maps are isomorphisms of vector spaces. The rest is a straightforward computation using the definitions

$$\eta_{[\sigma][\beta]} = \langle \tau_0(e_{[\sigma]})\tau_0(e_{[\beta]})\tau_0(e_{[1]}) \rangle_0^G$$

and

$$e_{[\alpha]} * e_{[\beta]} := \sum_{[\sigma], [\gamma]} \langle \tau_0(e_{[\alpha]})\tau_0(e_{[\beta]})\tau_0(e_{[\sigma]}) \rangle_0^G \eta^{[\sigma][\gamma]} e_{[\gamma]}.$$

□

These definitions are essentially the same as those given in [AGV], but they differ *a priori* from those of Chen and Ruan [CR1, Ru] for the “orbifold Poincaré pairing” and “orbifold cup product.” Nevertheless, it is easy to check that both the geometry and the final calculations are identical to those in [CR1, Ru]. For example, the three-fold multisectors $X_{[\gamma]}$ of [Ru, Defn 3.1.3] are, in our case, precisely the components $\overline{\mathcal{M}}_{0,3}(\mathcal{BG}, [\gamma])$.

We are interested now in the corresponding cohomological field theory (CohFT) and the large phase space (when $a_i > 0$).

Proposition 3.4. *The correlators $\langle \tau_{a_1}(e_{[\sigma_1]}) \dots \tau_{a_n}(e_{[\sigma_n]}) \rangle_g^G$ are related to the usual correlators $\langle \tau_{a_1} \dots \tau_{a_n} \rangle_g$ (corresponding to the case of $G = \{1\}$) by*

$$\langle \tau_{a_1}(e_{[\sigma_1]}) \dots \tau_{a_n}(e_{[\sigma_n]}) \rangle_g^G = \langle \tau_{a_i} \dots \tau_{a_n} \rangle_g \Omega_g^G(\gamma)$$

where

$$\Omega_g^G(\gamma) = \frac{|\mathcal{X}_g^G(\gamma)|}{|G|},$$

and

$$\mathcal{X}_g^G(\gamma) := \{(\alpha_1, \dots, \alpha_g, \beta_1, \dots, \beta_g, \sigma_1, \dots, \sigma_n) \mid \prod_{i=1}^g [\alpha_i, \beta_i] = \prod_{j=1}^n \sigma_j, \sigma_j \in [\gamma_j] \text{ for all } j\}.$$

Proof. The sublocus of $\overline{\mathcal{M}}_{g,n}(\mathcal{BG})$ where $\prod_{i=1}^n ev_i^*(e_{[\gamma_i]})$ is non-zero is $\overline{\mathcal{M}}_{g,n}(\mathcal{BG}, [\gamma_1], \dots, [\gamma_n])$. Proposition 2.1 shows that the degree of the forgetful map $\pi : \overline{\mathcal{M}}_{g,n}(\mathcal{BG}, [\gamma_1], \dots, [\gamma_n]) \longrightarrow \overline{\mathcal{M}}_{g,n}$ is exactly $\Omega_g^G(\gamma)$. The proof now follows from the projection formula, since ψ_i on $\overline{\mathcal{M}}_{g,n}(\mathcal{BG})$ is just the pullback $\pi^* \psi_i$ of the corresponding class on $\overline{\mathcal{M}}_{g,n}$. □

Lemma 3.5. *The numbers $\Omega_g^G(\gamma)$ depend only on the conjugacy classes $[\gamma_i]$, are independent of the ordering of the γ_i in γ , and satisfy the following relations:*

- (1) **Cutting trees:** For $g = g_1 + g_2$ and $I \amalg J = \{1, \dots, n\}$, let $\gamma_I = (\gamma_{i_1}, \dots, \gamma_{i_{|I|}})$ and $\gamma_J = (\gamma_{j_1}, \dots, \gamma_{j_{|J|}})$

$$\Omega_g^G(\gamma) = \Omega_{g_1}^G(\gamma_I, \zeta) \eta^{[\zeta][\xi]} \Omega_{g_2}^G(\xi, \gamma_J)$$

- (2) **Cutting loops:**

$$\Omega_g^G(\gamma) = \eta^{[\zeta][\xi]} \Omega_{g-1}^G(\zeta, \xi, \gamma)$$

- (3) **Forgetting tails:**

$$\Omega_g^G(\gamma) = \Omega_g^G(1, \gamma)$$

Proof. Independence of conjugacy class representative and of order are immediate from the definition, as is relation 3 (Forgetting tails). To prove relation 2, note that we may assume that $[\zeta] = [\xi^{-1}]$. Let $\mathcal{X}_g^G(\gamma)$ be as in Proposition 3.4 and let $\mathcal{Y}$ be the set

$$\mathcal{Y} := \{(\alpha'_1, \dots, \alpha'_{g-1}, \beta'_1, \dots, \beta'_{g-1}, \sigma'_1, \dots, \sigma'_n, \sigma'_{n+1}, \sigma'_{n+2}) \mid \prod_{i=1}^{g-1} [\alpha'_i, \beta'_i] = \prod_{j=1}^{n+2} \sigma'_j, \sigma'_j \in [\gamma_j] \text{ for } j \leq n, \sigma'_{n+2} \in [\sigma'^{-1}_{n+1}]\}.$$

Define a map $f : \mathcal{X}_g^G(\gamma) \longrightarrow \mathcal{Y}$, taking $(\alpha_1, \dots, \alpha_g, \beta_1, \dots, \beta_g, \sigma_1, \dots, \sigma_n)$ in $\mathcal{X}_g^G(\gamma)$ to $(\alpha_1, \dots, \alpha_{g-1}, \beta_1, \dots, \beta_{g-1}, \sigma_1, \dots, \sigma_n, \alpha_g \beta_g \alpha_g^{-1}, \beta_g^{-1})$ in $\mathcal{Y}$.

For a given conjugacy class $[\psi]$ in G , the map f , restricted to the subset of $\mathcal{X}_g^G(\gamma)$ where $\beta_g \in [\psi]$, takes $|C(\psi)|$ elements of $\mathcal{X}_g^G(\gamma)$ to one element of $\mathcal{Y}$. Moreover,

given $\zeta, \xi \in G$ such that $[\zeta^{-1}] = [\xi]$, if we let $\beta_g = \xi^{-1}$ and α_g be an element such that $\alpha_g \beta_g \alpha_g^{-1} = \zeta$, then it is clear that the map f is surjective. This shows that $\Omega_g^G(\gamma) = \sum_{[\zeta]} |C(\zeta)| \Omega_{g-1}^G(\gamma, \zeta, \zeta^{-1})$, which proves (2). The proof of (1) is a similar, straightforward argument. $\square$

Theorem 3.6. *Let $\Lambda_{g,n}^G : \mathcal{H}^{\otimes n} \longrightarrow H^*(\overline{\mathcal{M}}_{g,n})$ be defined as $\Lambda_{g,n}^G(e_{[\gamma_1]} \otimes \cdots \otimes e_{[\gamma_n]}) = \pi_*(ev_1^*(e_{[\gamma_1]}) \cdots ev_n^*(e_{[\gamma_n]})) = \Omega_g^G(\gamma)$. The collection $(\mathcal{H}, \eta, \Lambda_{g,n}^G, e_{[1]})$ is a CohFT with flat identity.*

Proof. It is straightforward to see that $\Lambda_{g,n}^G(e_{[\gamma_1]} \otimes \cdots \otimes e_{[\gamma_n]}) = \Omega_n^G(\gamma)$. The CohFT axioms follow immediately from Lemma 3.5. $\square$

The CohFT axioms also follow from the cutting and forgetting tails axioms that hold for the virtual fundamental class [CR2], which is trivial for $\overline{\mathcal{M}}_{g,n}(\mathcal{B}G)$. Alternatively, the axioms can be seen directly from the geometry by carefully accounting for ramification of π on the boundary and accounting for the degree of the morphism

$$\overline{\mathcal{M}}_{g,n}(\mathcal{B}G) \times_{\overline{\mathcal{M}}_{g,n}} (\overline{\mathcal{M}}_{g,n_1+1} \amalg \overline{\mathcal{M}}_{g_2,n_2+1}) \longrightarrow \overline{\mathcal{M}}_{g_1,n_1+1}(\mathcal{B}G) \amalg \overline{\mathcal{M}}_{g_2,n_2+1}(\mathcal{B}G)$$

as in [JKV, §4.2].

3.2. K-Theory. The orbifold K-theory of $\mathcal{B}G$ is simply the representation ring R_G , since a vector bundle on $\mathcal{B}G$ is a vector space with linear G -action. The Chern character (see [AR]) $ch : K_{orb}^* \otimes \mathbb{C} \longrightarrow H_{orb}^*(\mathcal{B}G, \mathbb{C}) = \mathcal{H}$ is easily seen to be the composite

$$R_G \otimes \mathbb{C} \xrightarrow{\chi} \text{Class}_{\mathbb{C}}(G) \cong \mathcal{H}$$

of the trace map χ and the obvious additive isomorphism from class functions to $\mathcal{H}$.

The trace map χ is a ring isomorphism, but as stated in Corollary 3.3, the orbifold product $*$ on $\mathcal{H}$ corresponds to convolution $\star$ of class functions rather than multiplication. Thus the Chern character is only an additive isomorphism.

3.3. Functoriality.

3.3.1. Morphisms.

It is interesting to note that K-theory has functoriality properties that H_{orb}^* does not enjoy. In particular, a homomorphism of groups $G \longrightarrow H$ gives a morphism $\mathcal{B}G \longrightarrow \mathcal{B}H$ which induces a ring homomorphism

$$K_{orb}^*(\mathcal{B}H) \longrightarrow K_{orb}^*(\mathcal{B}G),$$

corresponding to the obvious homomorphism of representation rings. But the center $Z\mathbb{C}[H]$ generally has no ring homomorphism to $Z\mathbb{C}[G]$.

3.3.2. Tensor products.

Tensor products arise in this theory in at least two ways.

- $\mathcal{B}(G \times H)$

For any two finite groups G and H , the classifying stack $\mathcal{B}(G \times H)$ splits up as a product

$$\mathcal{B}(G \times H) = \mathcal{B}G \times \mathcal{B}H,$$

and although the moduli stack of the product is not quite the product of the moduli stacks of the factors, it is easy to see that the corresponding CohFT

is the tensor product of the two components. That is, a straightforward check shows that if $\gamma = (\gamma_1, \dots, \gamma_n) \in G^n$ and $\sigma = (\sigma_1, \dots, \sigma_n) \in H^n$ then

$$\Omega_g^{G \times H}((\gamma_1, \sigma_1), \dots, (\gamma_n, \sigma_n)) = \Omega_g^G(\gamma) \Omega_g^H(\sigma).$$

• **$[X/G]$ with trivial G action**

If X is a smooth projective variety with trivial G action, then the quotient stack $[X/G]$ is isomorphic to the product $X \times \mathcal{B}G$.

Proposition 3.7. *For a smooth projective variety X with finite group G acting trivially, the CohFT arising from stable maps into the orbifold $[X/G]$ is simply the tensor product of the CohFT arising from stable maps into X and the CohFT arising from stable maps into $\mathcal{B}G$.*

Proof. This follows from the fact that the degree of the forgetful maps $\overline{\mathcal{M}}_{g,n}(X \times \mathcal{B}G, \llbracket \gamma_1 \rrbracket, \dots, \llbracket \gamma_n \rrbracket) \rightarrow \overline{\mathcal{M}}_{g,n}(X)$ and $\overline{\mathcal{M}}_{g,n}(\mathcal{B}G, \llbracket \gamma_1 \rrbracket, \dots, \llbracket \gamma_n \rrbracket) \rightarrow \overline{\mathcal{M}}_{g,n}$ are both equal to $\Omega_g(\gamma)$. $\square$

4. SEMISIMPLICITY AND VIRASORO ALGEBRAS

In this section, the summation convention is NOT used on any subscripts or superscripts α or α_i , although it is applied to all other variables.

4.1. Semisimple Frobenius algebras. Let V be any r -dimensional Frobenius algebra with multiplication $*$, metric η , and identity element $\mathbf{1}$. It is said to be a *semisimple Frobenius algebra* if there exists a *canonical basis* $\{f_\alpha\}_{\alpha=1}^r$ such that for all $\alpha_1, \alpha_2 = 1, \dots, r$,

$$(4.1) \quad f_{\alpha_1} * f_{\alpha_2} = \delta_{\alpha_1, \alpha_2} f_{\alpha_1}$$

and

$$(4.2) \quad \eta(f_{\alpha_1}, f_{\alpha_2}) = \delta_{\alpha_1, \alpha_2} \nu_{\alpha_1}$$

for some nonzero numbers ν_α . The identity element satisfies

$$\mathbf{1} = \sum_{\alpha=1}^r f_\alpha.$$

As discussed before, the Frobenius algebra $(\mathcal{H}, \eta, *, e_{\llbracket 1 \rrbracket})$ can be identified with $Z\mathbb{C}[G]$ and the latter is a semisimple Frobenius algebra with canonical basis given as follows.

Proposition 4.1. *Let $\{V_\alpha\}_{\alpha=1}^r$ be the set of irreducible representations of G and let χ_α denote the character of V_α . For all $\alpha = 1, \dots, r$, the elements*

$$f_\alpha := \frac{\dim V_\alpha}{|G|} \sum_{g \in G} \chi_\alpha(g^{-1}) g$$

form a basis of $Z\mathbb{C}[G]$ and satisfy equations (4.1) and (4.2), where for all $\alpha = 1, \dots, r$,

$$\nu_\alpha = \left(\frac{\dim V_\alpha}{|G|} \right)^2.$$

Proof. It is clear from the definition that the f_α lie in $Z\mathbb{C}[G]$. The fact that they satisfy (4.1) follows from [FH, §2.4]. That equation (4.2) holds for the given values of ν_α is a straightforward computation. $\square$

The results in the remainder of this section hold for any semi-simple Frobenius algebra, since any Frobenius algebra is a CohFT.

4.2. The Potential Function. We will now calculate the correlators for our theory in the canonical coordinates.

Proposition 4.2. *Let (g, n) be any stable pair, where $n \geq 1$. If $\alpha_i = \alpha$ for all $i = 1, \dots, n$ then for all $a_1, \dots, a_n \geq 0$, we have*

$$(4.3) \quad \langle \tau_{a_1}(f_\alpha) \cdots \tau_{a_n}(f_\alpha) \rangle_g^G = \nu_\alpha^{1-g} \langle \tau_{a_1} \cdots \tau_{a_n} \rangle_g;$$

otherwise, we have $\langle \tau_{a_1}(f_{\alpha_1}) \cdots \tau_{a_n}(f_{\alpha_n}) \rangle_g^G = 0$. Furthermore, when $n = 0$, we have $\langle \rangle_g^G = 0$.

Proof. Of course, $\langle \rangle_g^G = 0$ holds for dimensional reasons.

The proof for the rest of the proposition follows by degenerating to curves whose irreducible components are all three-pointed, genus-zero curves, where the proposition is easily verified, and then calculating the general correlators from the cutting axioms for CohFTs.

More explicitly, the correlator is simply

$$\langle \tau_{a_1}(f_{\alpha_1}) \cdots \tau_{a_n}(f_{\alpha_n}) \rangle_g^G = \Lambda_{g,n}^G(f_{\alpha_1} \otimes \cdots \otimes f_{\alpha_n}) \langle \tau_{a_1} \cdots \tau_{a_n} \rangle_g,$$

where $(\mathcal{H}, \eta, \Lambda^G)$ is our CohFT. The definition of $*$ gives

$$\begin{aligned} \Lambda_{0,3}^G(f_{\alpha_1} \otimes f_{\alpha_2} \otimes f_{\alpha_3}) &= \eta(f_{\alpha_1} * f_{\alpha_2}, f_{\alpha_3}) \\ &= \delta_{\alpha_1, \alpha_2} \delta_{\alpha_1, \alpha_3} \nu_{\alpha_1}. \end{aligned}$$

Now proceed by induction on the genus and number of marked points. Each application of the *cutting trees* axiom (for a 3-pointed, genus-zero vertex) leaves the genus unchanged, and reduces the number of marked points by one, but contributes nothing to the final result, since the node (cut edge) contributes the inverse metric—a factor of ν_α^{-1} —and the 3-pointed, genus-zero, irreducible component (vertex of the dual graph) contributes a factor of ν_α .

Each application of the *cutting loops* axiom increases the number of marked points by 2, reduces the genus by 1, and contributes the inverse of the metric—namely, ν_α^{-1} —to the final result. $\square$

The *large phase space potential* is defined by $\Phi^G(\mathbf{t}) = \sum_g \Phi_g^G(\mathbf{t}) \lambda^{2g-2}$ in $\lambda^{-2} \mathbb{C}[[\mathbf{t}, \lambda]]$, where $\Phi_g^G(\mathbf{t}) := \langle \exp(\boldsymbol{\tau} \cdot \mathbf{t}) \rangle_g^G$, where $\mathbf{t} \cdot \boldsymbol{\tau} = \sum_{a,m} t_a^m \tau_a(h_m)$, and where $\{h_m\}$ is any basis for $\mathcal{H}$. Let $Z^G := \exp(\Phi^G)$.

When $G = \{1\}$ is the trivial group, we denote by $\Phi := \Phi^{\{1\}}$, the potential of the Gromov-Witten invariants of a point. Similarly, we let $Z := Z^{\{1\}}$.

Proposition 4.3. *Let $\mathbf{u}$ be formal variables $\{u_a^\alpha\}$ for all integers $a \geq 0$ and $\alpha = 1, \dots, r$ associated to the canonical basis $\{f_\alpha\}$. For each $\alpha = 1, \dots, r$, let $\tilde{\mathbf{u}}^\alpha$ be formal variables $\{\tilde{u}_a^\alpha\}$, where $a \geq 0$ and $\tilde{u}_a^\alpha := (\nu_\alpha)^{\frac{1-a}{3}} u_a^\alpha$. Then we have*

$$(4.4) \quad \Phi^G(\mathbf{u}) = \sum_{\alpha=1}^r \Phi(\tilde{\mathbf{u}}^\alpha).$$

Proof. This follows from Proposition 4.2 and dimensional considerations. $\square$

4.3. Virasoro algebras. For each $\alpha = 1, \dots, r$ and $n \geq -1$, let

$$\begin{aligned} L_n^{(\alpha)} &:= -\frac{(2n+3)!!}{2^{n+1}} \frac{\partial}{\partial \tilde{u}_{n+1}^\alpha} + \sum_{i=0}^{\infty} \frac{(2i+2n+1)!!}{(2i-1)!!2^{n+1}} \tilde{u}_i^\alpha \frac{\partial}{\partial \tilde{u}_{i+n}^\alpha} \\ &+ \frac{\lambda^2}{2} \sum_{i=0}^{n-1} \frac{(2i+1)!!(2n-(2i+1))!!}{2^{n+1}} \frac{\partial^2}{\partial \tilde{u}_i^\alpha \partial \tilde{u}_{n-1-i}^\alpha} \\ &+ \delta_{n,-1} \frac{\lambda^{-2}}{2} \tilde{u}_0^\alpha \tilde{u}_0^\alpha + \delta_{n,0} \frac{1}{16}, \end{aligned}$$

where $(2n-1)!! := 1 \cdot 3 \cdot 5 \dots (2n-1)$. These operators satisfy

$$[L_m^{(\alpha)}, L_n^{(\beta)}] = (m-n)L_{m+n}^{(\alpha)} \delta_{\alpha,\beta}$$

for all $m, n \geq -1$ and $\alpha, \beta \in \{1, \dots, r\}$, forming r commuting copies of “half” of the Virasoro algebra.

Moreover, if $\{b_m\}$ is any basis for $\mathcal{H}$, where $m = 0, \dots, r-1$ such that $b_0 = \mathbf{1}$, and if $\mathbf{t} = \{t_a^m\}$ are the associated formal parameters, then there are also operators for all $n \geq -1$ given by

$$\begin{aligned} L_n &:= -\frac{(2n+3)!!}{2^{n+1}} \frac{\partial}{\partial t_{n+1}^0} + \sum_{i=0}^{\infty} \frac{(2i+2n+1)!!}{(2i-1)!!2^{n+1}} \left(\sum_m t_i^m \frac{\partial}{\partial t_{i+n}^m} \right) \\ &+ \frac{\lambda^2}{2} \sum_{i=0}^{n-1} \frac{(2i+1)!!(2n-(2i+1))!!}{2^{n+1}} \left(\sum_{m_1, m_2} \eta^{m_1 m_2} \frac{\partial^2}{\partial t_i^{m_1} \partial t_{n-1-i}^{m_2}} \right) \\ &+ \delta_{n,-1} \frac{\lambda^{-2}}{2} \left(\sum_{m_1, m_2} \eta_{m_1 m_2} t_0^{m_1} t_0^{m_2} \right) + \delta_{n,0} \frac{r}{16}, \end{aligned}$$

satisfying $[L_k, L_n] = (k-n)L_{k+n}$ for any $k, n \geq -1$.

Proposition 4.4. For all $\alpha \in \{1, \dots, r\}$ and $n \geq -1$,

$$(4.5) \quad L_n^{(\alpha)} Z^G = 0.$$

These equations completely determine Z^G . Furthermore, for all $n \geq -1$,

$$(4.6) \quad L_n Z^G = 0.$$

Proof. Equation (4.5) follows from Proposition 4.3 and the Kontsevich-Witten theorem [Ko, Wi] for the case of $G = \{\mathbf{1}\}$. Equation (4.6) follows from (4.5) and the identity

$$L_m = \sum_{\alpha=1}^r (\nu_\alpha)^{-\frac{m}{3}} L_m^{(\alpha)}.$$

□

Remark 4.5. Equation 4.6 is a verification of the Virasoro conjecture for $\mathcal{BG}$ [EHX] and can also be regarded as an example of [Gi].

4.4. KdV hierarchies. Let $\langle\langle A \rangle\rangle_g^G := \langle\langle A \exp(\mathbf{t} \cdot \boldsymbol{\tau}) \rangle\rangle_g^G$ and $\langle\langle A \rangle\rangle^G := \sum_g \langle\langle A \exp(\mathbf{t} \cdot \boldsymbol{\tau}) \rangle\rangle_g^G \lambda^{2g-2}$. The superscript G will be suppressed when $G = \{\mathbf{1}\}$, the trivial group.

Proposition 4.6. *Let $\{e_1, \dots, e_m\}$ be any basis for $\mathcal{H}$. For all v in $\mathcal{H}$ and $a \geq 0$, the following equation holds:*

$$(4.7) \quad \begin{aligned} & (2a+1)\lambda^{-2} \langle\langle \tau_a(v) \tau_0(e_{m_1}) \tau_0(e_{m_2}) \rangle\rangle^G \eta^{m_1 m_2} = \\ & \langle\langle \tau_{a-1}(v) \tau_0(e_{m_1}) \rangle\rangle^G \eta^{m_1 m_2} \langle\langle \tau_0(e_{m_2}) \tau_0(e_{m_3}) \tau_0(e_{m_4}) \rangle\rangle^G \eta^{m_3 m_4} + \\ & 2 \langle\langle \tau_{a-1}(v) \tau_0(e_{m_1}) \tau_0(e_{m_3}) \rangle\rangle^G \eta^{m_1 m_2} \eta^{m_3 m_4} \langle\langle \tau_0(e_{m_2}), \tau_0(e_{m_4}) \rangle\rangle^G \eta^{m_3 m_4} + \\ & \frac{1}{4} \langle\langle \tau_{a-1}(v) \tau_0(e_{m_1}) \tau_0(e_{m_2}) \tau_0(e_{m_3}) \tau_0(e_{m_4}) \rangle\rangle^G \eta^{m_1 m_2} \eta^{m_3 m_4}. \end{aligned}$$

Equation (4.7) and the fact that $L_{-1}^{(\alpha)} Z^G = 0$ for all $\alpha = 1, \dots, r$ completely determine Φ^G .

Proof. When $G = \{\mathbf{1}\}$, the trivial group, equation (4.7) reduces to

$$(4.8) \quad \begin{aligned} & (2a+1)\lambda^{-2} \langle\langle \tau_a \tau_0 \tau_0 \rangle\rangle = \\ & \langle\langle \tau_{a-1} \tau_0 \rangle\rangle \langle\langle \tau_0 \tau_0 \tau_0 \rangle\rangle + 2 \langle\langle \tau_{a-1} \tau_0 \tau_0 \rangle\rangle \langle\langle \tau_0, \tau_0 \rangle\rangle + \frac{1}{4} \langle\langle \tau_{a-1} \tau_0 \tau_0 \tau_0 \tau_0 \rangle\rangle. \end{aligned}$$

This equation is the Kontsevich-Witten theorem [Ko, Wi]. Witten also showed that this equation together with $L_{-1} Z = 0$ completely determines Φ .

To prove the formula for general G , choose a canonical basis $\{f_\alpha\}_{\alpha=1}^r$ and let $v = f_\alpha$. Consider the terms of equation (4.7) proportional to λ^{2g-4} . By Proposition (4.2), one obtains the terms of equation (4.7) proportional to λ^{2g-4} up to an overall scalar factor. $\square$

Remark 4.7. Equation 4.7 is a simultaneous solution of r commuting KdV hierarchies with time parameters $\tilde{u}_a^\alpha$ for $a \geq 0$ and $\alpha = 1, \dots, r$.

REFERENCES

- ACV. D. Abramovich, A. Corti, and A. Vistoli, *Twisted bundles and admissible covers*, math.AG/0106211.
- AGV. D. Abramovich, T. Graber, and A. Vistoli, *Algebraic orbifold quantum products*, math.AG/0112004. To appear in this volume.
- AV. D. Abramovich, A. Vistoli, *Compactifying the space of stable maps*, J. Amer. Math. Soc. **15**, no. 1 (2001), 27–75. math.AG/9908167.
- AR. A. Adem, Y. Ruan, *Twisted orbifold K-theory*, math.AT/0107168 v1.
- CR1. W. Chen, Y. Ruan, *A new cohomology theory for orbifold*, math.AG/0004129.
- CR2. ———, *Orbifold Gromov-Witten theory*, math.AG/0103156.
- EHX. T. Eguchi, K. Hori, Ch.-Sh. Xiong, *Quantum cohomology and the Virasoro algebra*, Phys. Lett. **B402** (1997), 71–80.
- FH. W. Fulton, J. Harris, *Representation theory, A first course*, Springer-Verlag, 1991.
- Gi. A. Givental, *Gromov-Witten invariants and quantization of quadratic Hamiltonians*, math.AG/0108100.
- JKV. T. J. Jarvis, T. Kimura, and A. Vaintrob, *Moduli spaces of higher spin curves and integrable hierarchies*, Compositio Math. **126** (2001) 157–212, math.AG/9905034.
- Ko. M. Kontsevich, *Intersection theory on the moduli space of curves and the matrix Airy function*, Commun. Math. Phys. **147** (1992), 1–23.
- Ru. Y. Ruan, *Discrete torsion and twisted orbifold cohomology*, math.AG/0005299.
- Wi. E. Witten, *Two-dimensional gravity and intersection theory on moduli space*, Surveys in Diff. Geom. **1** (1991), 243–310.

DEPARTMENT OF MATHEMATICS, BRIGHAM YOUNG UNIVERSITY, PROVO, UT 84602, USA

E-mail address: `jarvis@math.byu.edu`

DEPARTMENT OF MATHEMATICS, 111 CUMMINGTON STREET, BOSTON UNIVERSITY, BOSTON,
MA 02215, USA

E-mail address: `kimura@math.bu.edu`